

Advanced Wound Care



Antibacterial **MEDIHONEY**[®] *Wound Gel*

MediHoney[®] Antibacterial Wound Gel consists of Medical Grade Manuka Honey combined with gelling agents derived from natural sources.

80%

**ANTIBACTERIAL
MANUKA HONEY***

Features and Benefits

- Made with Medical Grade Manuka Honey
- Delivers broad spectrum antimicrobial activity¹
- Helps to reduce malodour²
- MediHoney aids and supports autolytic debridement and a moist wound healing environment in acute and chronic wounds and burns³⁻⁶
- MediHoney has a low pH of 3.5-4.5
- Lowering pH has been associated with wound healing benefits
- Has been safely used on patients of all ages
- Dressing change frequency will depend on the condition of the patient as well as the level of wound exudate

MEDIHONEY® Wound Gel


Antibacterial Dressing

MediHoney® Wound Gel Antibacterial Dressing

Reference	Description	Units/Box	PIP Code	NHS Code
391	10g	20/box	314-1207	ELZ013
395	20g	5/box	314-1215	ELZ507



Peer-review study to show the % of MediHoney® required to kill specific organisms⁷



Indications

- Acute and chronic wounds
- Infected and malodorous wounds
- Necrotic and sloughy wounds
- Pressure ulcers
- Surgical, donor sites, and recipient graft sites
- Leg ulcers (venous, arterial and mixed aetiology ulcers) and diabetic foot ulcers
- Superficial burns
- Superficial wounds such as cuts, scratches, and abrasions

Contraindications

- Third degree burns
- Patients with a known sensitivity to honey
- To control heavy bleeding

References: 1. Maddocks S, Lopez M, Rowlands R, Cooper R. Manuka honey inhibits the development of Streptococcus pyogenes biofilms and causes reduced expression of two fibronectin binding proteins. Microbiology. 2012;158:781-790. 2. Robson V, Yorke J, Sen R, Lowe D, Rogers S. Randomised controlled feasibility trial on the use of medical grade honey following microvascular free tissue transfer to reduce the incidence of wound infection. The British J Oral & Maxillofacial Surgery. 2012;50(4):321-327. 3. Robson, V., Dodd, S and Thomas, S. Standardized antibacterial honey (MediHoney®) with standard therapy in wound care: randomized clinical trial. Journal of Advanced Nursing. 2009; 565-575. 4. Bateman S, Graham T. The Use of MediHoney® Wound Gel on surgical wounds post-CABG. WOUNDS UK. 2007;(3):76 – 83. 5. Cadogan, J. The use of honey to treat an ulcer on the heel of a person with diabetes. The Diabetic Foot Journal. 2008 11. (1): 43-45. 6. In house data. 7. George N, Cutting K. Antibacterial Honey (Medihoney™): in-vitro Activity against Clinical Isolates of MRSA, VRE, and Other Multiresistant Gram-negative Organisms Including Pseudomonas Aeruginosa. Wounds. 2007; 19(9):231-236.

Availability of these products might vary from a given country or region to another, as a result of specific local regulatory approval or clearance requirements for sale in such country or region.

- Non contractual document. The manufacturer reserves the right, without prior notice, to modify the products in order to improve their quality.
- Consult product labels and inserts for any indication, contraindications, hazards, warnings, precautions, and instructions for use.

Additional information for EMEA Customers only:

Products mentioned in this document are CE class I, IIa, IIb or III devices. Contact Integra should you need any additional information on devices classification. All the medical devices mentioned on this document are CE marked according to European council directive 93/42/EEC on medical devices and its relatives, unless specifically identified as "NOT CE MARKED".

For more information or to place an order, please contact:

Customer Service

United Kingdom +44 (0) 1264 312 725 ■ +44 (0) 1264 312 821 fax
integrallife.eu



Manufacturer:



Derma Sciences, Inc.
 104 Shorting Road
 Toronto, Ontario M1S 3S4 ■ Canada



Scanlan Group B.V.
 Postbus 75664
 Schiphol Triport 1118 ZS
 The Netherlands
 Int: +31 20 653-0553